


3D Transformations

CMSC 435/634


Transformation

Webster: The operation of changing one configuration or expression into another in accordance with a mathematical rule



Using Transformation

- ▶ Points on object represented as vector offset from origin
- ▶ Transform is a vector to vector function
 - ▶ $\vec{p}' = f(\vec{p})$
- ▶ Relativity:
 - ▶ From \vec{p}' point of view, object is transformed
 - ▶ From \vec{p} point of view, coordinate system changes
- ▶ Inverse transform, $\vec{p} = f^{-1}(\vec{p}')$




Composing Transforms

- ▶ Order matters

- ▶ $R(T(\vec{p})) = R \circ T(\vec{p})$


- ▶ $T(R(\vec{p})) = T \circ R(\vec{p})$



Inverting Composed Transforms

- ▶ Reverse order

- ▶ $(R \circ T)^{-1}(\vec{p}') = T^{-1}(R^{-1}(\vec{p}'))$
- ▶ $(T \circ R)^{-1}(\vec{p}') = R^{-1}(T^{-1}(\vec{p}'))$




Translation

▶ $\vec{p}' = \vec{p} + \vec{t}$

▶
$$\begin{bmatrix} p'^x \\ p'^y \\ p'^z \end{bmatrix} = \begin{bmatrix} p^x \\ p^y \\ p^z \end{bmatrix} + \begin{bmatrix} t^x \\ t^y \\ t^z \end{bmatrix} = \begin{bmatrix} p^x + t^x \\ p^y + t^y \\ p^z + t^z \end{bmatrix}$$

▶ \vec{t} says where \vec{p} -space origin ends up ($\vec{p}' = \vec{0} + \vec{t}$)

▶ Composition: $\vec{p}' = (\vec{p} + \vec{t}_0) + \vec{t}_1 = \vec{p} + (\vec{t}_0 + \vec{t}_1)$



Linear Transforms

$$\begin{bmatrix} p'^x \\ p'^y \\ p'^z \end{bmatrix} = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} \begin{bmatrix} p^x \\ p^y \\ p^z \end{bmatrix}$$

- Matrix says where \vec{p} -space axes end up

$$\begin{bmatrix} a \\ d \\ g \end{bmatrix} = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} b \\ e \\ h \end{bmatrix} = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$$


$$\begin{bmatrix} c \\ f \\ i \end{bmatrix} = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

- Composition: $\vec{p}' = M (N \vec{p}) = (M N) \vec{p}$

Common case: Scaling

$$\blacktriangleright \begin{bmatrix} p'^x \\ p'^y \\ p'^z \end{bmatrix} = \begin{bmatrix} s_x p^x \\ s_y p^y \\ s_z p^z \end{bmatrix} = \begin{bmatrix} s_x & 0 & 0 \\ 0 & s_y & 0 \\ 0 & 0 & s_z \end{bmatrix} \begin{bmatrix} p^x \\ p^y \\ p^z \end{bmatrix}$$

$$\blacktriangleright \text{Inverse: } \begin{bmatrix} 1/s_x & 0 & 0 \\ 0 & 1/s_y & 0 \\ 0 & 0 & 1/s_z \end{bmatrix}$$




Common case: Reflection

- ▶ Negative scaling

$$\begin{bmatrix} p'^x \\ p'^y \\ p'^z \end{bmatrix} = \begin{bmatrix} -p^x \\ p^y \\ p^z \end{bmatrix} = \begin{bmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} p^x \\ p^y \\ p^z \end{bmatrix}$$




Common case: Rotation



- ▶ Orthogonal, so $M^{-1} = M^T$

- ▶ Rotate around Z: $\vec{p}' = \begin{bmatrix} \cos \theta & -\sin \theta & 0 \\ \sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{bmatrix} \vec{p}$


Common case: Rotation



- ▶ Orthogonal, so $M^{-1} = M^T$

- ▶ Rotate around X: $\vec{p}' = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \cos \theta & -\sin \theta \\ 0 & \sin \theta & \cos \theta \end{bmatrix} \vec{p}$

Common case: Rotation



- ▶ Orthogonal, so $M^{-1} = M^T$

- ▶ Rotate around Y: $\vec{p}' = \begin{bmatrix} \cos \theta & 0 & \sin \theta \\ 0 & 1 & 0 \\ -\sin \theta & 0 & \cos \theta \end{bmatrix} \vec{p}$

Composing Transforms

- ▶ Scale by s along axis \vec{a}
 - ▶ Rotate to align \vec{a} with Z
 - ▶ Scale along Z
 - ▶ Rotate back

Rotate by α around X into XZ plane

- ▶ Projection of \vec{a} onto YZ: $\begin{bmatrix} 0 \\ a^y \\ a^z \end{bmatrix}$
- ▶ length $d = \sqrt{(a^y)^2 + (a^z)^2}$
- ▶ So $\cos \alpha = a^z/d$, $\sin \alpha = a^y/d$
- ▶ $R_X = \begin{bmatrix} 1 & 0 & 0 \\ 0 & a^z/d & -a^y/d \\ 0 & a^y/d & a^z/d \end{bmatrix}$
- ▶ Result $\vec{a}' = \begin{bmatrix} a^x \\ 0 \\ d \end{bmatrix}$

Rotate by β around Y to Z axis

▶ $\vec{a}' = \begin{bmatrix} a^x \\ 0 \\ d \end{bmatrix}$

▶ length = 1

▶ So $\cos \beta = d$, $\sin \beta = a^x$

▶ $R_Y = \begin{bmatrix} d & 0 & -a^x \\ 0 & 1 & 0 \\ a^x & 0 & d \end{bmatrix}$

▶ Result $\vec{a}'' = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$

Composing Transforms

- ▶ Scale by s along Z : $S_Z = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & s \end{bmatrix}$
- ▶ Scale by s along axis \vec{a}
 - ▶ Rotate to align \vec{a} with Z
 - ▶ Scale along Z
 - ▶ Rotate back
 - ▶ $\vec{p}' = R_X^{-1} R_Y^{-1} S_Z R_Y R_X \vec{p}$

Affine Transforms

- ▶ Affine = Linear + Translation
- ▶ Composition? $A (B \vec{p} + \vec{t}_0) + \vec{t}_1 = A B \vec{p} + A \vec{t}_0 + \vec{t}_1$
- ▶ Yuck!

Homogeneous Coordinates

- ▶ Add a '1' to each point

$$\begin{bmatrix} p'^x \\ p'^y \\ p'^z \\ 1 \end{bmatrix} = \left[\begin{array}{ccc|c} a & b & c & t^x \\ d & e & f & t^y \\ g & h & i & t^z \\ \hline 0 & 0 & 0 & 1 \end{array} \right] \begin{bmatrix} p^x \\ p^y \\ p^z \\ 1 \end{bmatrix}$$

- ▶ $\vec{p}'^x = (a p^x + b p^y + c p^z) + t^x$
- ▶ $\vec{p}'^y = (d p^x + e p^y + f p^z) + t^y$
- ▶ $\vec{p}'^z = (g p^x + h p^y + i p^z) + t^z$
- ▶ $1 = (0p^x + 0p^y + 0p^z) + 1$

Homogeneous Coordinates

$$\begin{matrix} \blacktriangleright \end{matrix} \begin{bmatrix} p'^x \\ p'^y \\ p'^z \\ 1 \end{bmatrix} = \begin{bmatrix} a & b & c & | & t^x \\ d & e & f & | & t^y \\ g & h & i & | & t^z \\ \hline 0 & 0 & 0 & | & 1 \end{bmatrix} \begin{bmatrix} p^x \\ p^y \\ p^z \\ 1 \end{bmatrix}$$

- ▶ $\vec{p}' = [\vec{x} \ \vec{y} \ \vec{z} \ | \ \vec{t}] \vec{p}$
 - ▶ \vec{t} says where the \vec{p} -space origin ends up
 - ▶ $\vec{x}, \vec{y}, \vec{z}$ say where the \vec{p} -space axes end up
- ▶ Composition: Just matrix multiplies!

Composing Transforms

- ▶ Rotate by θ about line between \vec{p}_0 and \vec{p}_1 :
 - ▶ Translate \vec{p}_0 to origin
 - ▶ Rotate to align $\vec{p}_1 - \vec{p}_0$ with Z
 - ▶ Rotate by θ around Z
 - ▶ Undo $\vec{p}_1 - \vec{p}_0$ rotation
 - ▶ Undo translation
- ▶ $T^{-1}R_X^{-1}R_Y^{-1}R_Z(\theta)R_YR_XT$

Vectors

- ▶ Transform by *Jacobian Matrix*
- ▶ Matrix of partial derivatives

$$\begin{bmatrix} \vec{p}'^x \\ \vec{p}'^y \\ \vec{p}'^z \end{bmatrix} = \begin{bmatrix} a p^x + b p^y + c p^z + t^x \\ d p^x + e p^y + f p^z + t^y \\ g p^x + h p^y + i p^z + t^z \end{bmatrix}$$

- ▶ $J = \begin{bmatrix} \partial p'^x / \partial p^x a & \partial p'^x / \partial p^y b & \partial p'^x / \partial p^z c \\ \partial p'^y / \partial p^x d & \partial p'^y / \partial p^y e & \partial p'^y / \partial p^z f \\ \partial p'^z / \partial p^x g & \partial p'^z / \partial p^y h & \partial p'^z / \partial p^z i \end{bmatrix}$
- ▶ *Upper-left 3x3*

Normals

- ▶ Normal should remain perpendicular to tangent vector

- ▶ $\vec{n} \cdot \vec{v} = \vec{n}' \cdot \vec{v}' = 0$

- ▶
$$\begin{bmatrix} n_x & n_y & n_z \end{bmatrix} J^{-1} J \begin{bmatrix} v^x \\ v^y \\ v^z \end{bmatrix} = \begin{bmatrix} v'^x \\ v'^y \\ v'^z \end{bmatrix} = 0$$

- ▶ $\vec{n}' = \vec{n} J^{-1}$

- ▶ Multiply by inverse on right

- ▶ OR multiply *column* normal by inverse transpose

Nesting

- ▶ Room
 - ▶ Desk
 - ▶ Student
 - ▶ Book
 - ▶ Notebook
 - ▶ Desk
 - ▶ Student
 - ▶ Notebook
 - ▶ Table
 - ▶ Laptop
 - ▶ Blackboard
 - ▶ Chalk
 - ▶ Chalk
 - ▶ Eraser

Matrix Stack

- ▶ Remember transformation, return to it later
- ▶ Push a copy, modify the copy, pop
 - ▶ `RiBeginTransform()/RiEndTransform()`
 - ▶ `glPushMatrix()/glPopMatrix()`
- ▶ Keep matrix and update matrix and inverse
- ▶ Push and pop both