

Illumination

CMSC 435/634

Illumination

Local Illumination

Interpolation

Illumination

Local Illumination

Interpolation

Illumination

- ▶ Effect of light on objects
- ▶ Mostly look just at intensity
 - ▶ Apply to each color channel independently
- ▶ Good for most objects
 - ▶ Not fluorescent
 - ▶ Not phosphorescent

Local vs. Global

- ▶ Local
 - ▶ Light sources shining directly on object
- ▶ Global
 - ▶ Lights bouncing from objects onto other objects
 - ▶ Ambient Illumination
 - ▶ Approximate global illumination as constant color
 - ▶ Typically $\sim 1\%$ of direct illumination

Illumination

Local Illumination

BRDF

Rendering Equation


Models

Interpolation

BRDF


Bidirectional Reflectance Distribution Function

How much light reflects from L_i to L_o




Physically Plausible BRDF

- ▶ Positive
- ▶ Reciprocity
 - ▶ Same light from L_i to L_o as from L_o to L_i
- ▶ Conservation of Energy
 - ▶ Don't reflect more energy than comes in



Plotting BRDFs

- ▶ Polar plot of reflectance strength
 - ▶ For **one** view direction, showing light directions
 - ▶ For **one** light direction, showing view directions
- ▶ Reciprocity – same if you swap view and light



Rendering Equation

Integral of all Incoming Light

$$L_o(\hat{v}) = \int_{\Omega(\hat{n})} L_i(\hat{l}) f_r(\hat{v}, \hat{l}) \hat{n} \cdot \hat{l} d\omega(\hat{l})$$

Parts of this equation:

$L_o(\hat{v})$	outgoing light in direction \hat{v}
$\Omega(\hat{n})$	hemisphere above \hat{n} that can see this point
$L_i(\hat{l})$	incoming light from direction \hat{l}
$f_r(\hat{v}, \hat{l})$	BRDF from \hat{l} to \hat{v}
$\hat{n} \cdot \hat{l} d\omega(\hat{l})$	projection of differential solid angle onto surface

Rendering Equation for Point Lights

Sum for Each Light


$$L_o(\hat{v}) = \sum_i L_i f_r(\hat{v}, \hat{l}_i) \hat{n} \cdot \hat{l}_i$$

Parts of this equation:

- $L_o(\hat{v})$ outgoing light in direction \hat{v}
- i lights that can see this point (where $\hat{n} \cdot \hat{l}_i > 0$)
- \hat{l}_i light direction to light i
- L_i incoming light for light i
- $f_r(\hat{v}, \hat{l}_i)$ BRDF from \hat{l}_i to \hat{v}

Results

- ▶ Integrating full environment
- ▶ Light at one point, black elsewhere



Decomposing BRDFs

- ▶ Decompose BRDF into convenient parts
- ▶ Typical breakdown:
 - ▶ Diffuse (view independent)
 - ▶ Specular (view dependent near reflection)
 - ▶ Others less common, often ignored (e.g. retro reflection)




$$L_o(\hat{\mathbf{v}}) = \sum_i L_i \left(f_d(\hat{\mathbf{v}}, \hat{\mathbf{l}}_i) + f_s(\hat{\mathbf{v}}, \hat{\mathbf{l}}_i) \right) \hat{\mathbf{n}} \cdot \hat{\mathbf{l}}_i$$

$$L_o(\hat{\mathbf{v}}) = \sum_i L_i f_d(\hat{\mathbf{v}}, \hat{\mathbf{l}}_i) \hat{\mathbf{n}} \cdot \hat{\mathbf{l}}_i + \sum_i L_i f_s(\hat{\mathbf{v}}, \hat{\mathbf{l}}_i) \hat{\mathbf{n}} \cdot \hat{\mathbf{l}}_i$$


Important directions

\hat{n} : Unit surface normal




Important directions

\hat{v} : Unit vector from surface toward viewer




Important directions

\hat{i} : Unit vector from surface toward light




Important directions

$\hat{R}_v = 2\hat{n}(\hat{n} \cdot \hat{v}) - \hat{v}$: Direction of mirror reflection of view




Important directions

$\hat{R}_l = 2\hat{n}(\hat{n} \cdot \hat{l}) - \hat{l}$: Direction of mirror reflection of light



Important directions

$\hat{h} = (\hat{v} + \hat{l}) / |\hat{v} + \hat{l}|$: Normal direction that would reflect \hat{v} to \hat{l}




Diffuse

- ▶ Also called Lambertian or Matte
- ▶ Total reflectance: $\sum_i L_i Kd \hat{n} \cdot \hat{l}_i$
- ▶ BRDF: Kd



Phong

- ▶ Strongest where \hat{R}_l lines up with \hat{v} or \hat{R}_v lines up with \hat{l}
- ▶ Total reflectance: $\sum_i L_i K_s (\hat{R}_v \cdot \hat{l}_i)^e$
- ▶ Physically plausible version: $\sum_i L_i K_s (\hat{R}_v \cdot \hat{l}_i)^e \hat{n} \cdot \hat{l}$
 - ▶ With energy-conserving K_s



Specular Microfacets

- ▶ Imagine random mirrored *microfacets*



- ▶ Normal Distribution Function (NDF)
 - ▶ Probability facet has normal \hat{h}
 - ▶ Only facets to reflect \hat{l} to \hat{v}
- ▶ Proportion of light or view blocked (geometry term)
 - ▶ Blocked light = *shadowing*
 - ▶ Blocked view = *masking*
- ▶ Fresnel term
 - ▶ Reflection from non-metals is stronger at glancing angles

Cook-Torrance

- ▶ Beckmann Distribution = Gaussian distribution of slope
- ▶ Shadow/Mask based on symmetric V-shaped microfacets
- ▶ BRDF: $D(\hat{n}, \hat{h}) \frac{G(\hat{n}, \hat{v}, \hat{l})}{4 \hat{n} \cdot \hat{v} \hat{n} \cdot \hat{l}} F(\hat{v}, \hat{l})$,
- ▶ Total reflectance: $\sum_i L_i K_s D(\hat{n}, \hat{h}_i) \frac{G(\hat{n}, \hat{v}, \hat{l}_i)}{4 \hat{n} \cdot \hat{v} \hat{n} \cdot \hat{l}_i} F(\hat{v}, \hat{l}_i) \hat{n} \cdot \hat{l}$



Blinn-Phong

- ▶ Alternate formulation for Phong, similar behavior
- ▶ Strongest where \hat{h} lines up with \hat{n}
 - ▶ Function of \hat{h} , behaves like NDF
- ▶ Total reflectance (original form): $\sum_i L_i K_s (\hat{n} \cdot \hat{h}_i)^e$
- ▶ As NDF: $D(\hat{n}, \hat{h}_i) = \frac{e+2}{2\pi} (\hat{n} \cdot \hat{h}_i)^e$



Illumination

Local Illumination

Interpolation

When to Compute

- ▶ *Gouraud Shading* = Compute per-vertex & interpolate
 - ▶ Lose sharp highlights
 - ▶ Subject to *Mach banding*
- ▶ *Phong Shading* = Interpolate normals & compute per-pixel



Gouraud



Phong

Phong Shading

- ▶ Phong shading can refer to lighting model **or** interpolation
- ▶ To save confusion:
 - ▶ *Phong lighting*
 - ▶ *Phong interpolation*